

Measurement of the branching fractions for $\psi(2S) \rightarrow e^+e^-$ and $\psi(2S) \rightarrow \mu^+\mu^-$

- B. Aubert,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ P. Robbe,¹ V. Tisserand,¹ A. Palano,² A. Pompili,² G. P. Chen,³ J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ P. L. Reinertsen,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ A. R. Clark,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ J. F. Kral,⁵ C. LeClerc,⁵ M. E. Levi,⁵ T. Liu,⁵ G. Lynch,⁵ P. J. Oddone,⁵ A. Perazzo,⁵ M. Pripstein,⁵ N. A. Roe,⁵ A. Romosan,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ P. G. Bright-Thomas,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ D. J. Knowles,⁶ S. W. O'Neale,⁶ R. C. Penny,⁶ A. T. Watson,⁶ N. K. Watson,⁶ T. Deppermann,⁷ K. Goetzen,⁷ H. Koch,⁷ M. Kunze,⁷ B. Lewandowski,⁷ K. Peters,⁷ H. Schmuecker,⁷ M. Steinke,⁷ J. C. Andress,⁸ N. R. Barlow,⁸ W. Bhimji,⁸ N. Chevalier,⁸ P. J. Clark,⁸ W. N. Cottingham,⁸ N. De Groot,⁸ N. Dyce,⁸ B. Foster,⁸ J. D. McFall,⁸ D. Wallom,⁸ F. F. Wilson,⁸ K. Abe,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ S. Jolly,¹⁰ A. K. McKemey,¹⁰ J. Tinslay,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ D. A. Bukin,¹¹ A. R. Buzykaev,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ A. A. Korol,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ A. A. Salnikov,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ V. I. Telnov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² A. J. Lankford,¹² M. Mandelkern,¹² S. McMahan,¹² D. P. Stoker,¹² A. Ahsan,¹³ K. Arisaka,¹³ C. Buchanan,¹³ S. Chun,¹³ J. G. Branson,¹⁴ D. B. MacFarlane,¹⁴ S. Prell,¹⁴ Sh. Rahatlou,¹⁴ G. Raven,¹⁴ V. Sharma,¹⁴ C. Campagnari,¹⁵ B. Dahmes,¹⁵ P. A. Hart,¹⁵ N. Kuznetsova,¹⁵ S. L. Levy,¹⁵ O. Long,¹⁵ A. Lu,¹⁵ J. D. Richman,¹⁵ W. Verkerke,¹⁵ M. Witherell,¹⁵ S. Yellin,¹⁵ J. Beringer,¹⁶ D. E. Dorfan,¹⁶ A. M. Eisner,¹⁶ A. A. Grillo,¹⁶ M. Grothe,¹⁶ C. A. Heusch,¹⁶ R. P. Johnson,¹⁶ W. S. Lockman,¹⁶ T. Pulliam,¹⁶ H. Sadrozinski,¹⁶ T. Schalk,¹⁶ R. E. Schmitz,¹⁶ B. A. Schumm,¹⁶ A. Seiden,¹⁶ M. Turri,¹⁶ W. Walkowiak,¹⁶ D. C. Williams,¹⁶ M. G. Wilson,¹⁶ E. Chen,¹⁷ G. P. Dubois-Felsmann,¹⁷ A. Dvoretzki,¹⁷ D. G. Hitlin,¹⁷ S. Metzler,¹⁷ J. Oyang,¹⁷ F. C. Porter,¹⁷ A. Ryd,¹⁷ A. Samuel,¹⁷ M. Weaver,¹⁷ S. Yang,¹⁷ R. Y. Zhu,¹⁷ S. Devmal,¹⁸ T. L. Geld,¹⁸ S. Jayatilake,¹⁸ G. Mancinelli,¹⁸ B. T. Meadows,¹⁸ M. D. Sokoloff,¹⁸ T. Barillari,¹⁹ P. Bloom,¹⁹ M. O. Dima,¹⁹ S. Fahey,¹⁹ W. T. Ford,¹⁹ D. R. Johnson,¹⁹ U. Nauenberg,¹⁹ A. Olivas,¹⁹ P. Rankin,¹⁹ J. Roy,¹⁹ S. Sen,¹⁹ J. G. Smith,¹⁹ W. C. van Hoek,¹⁹ D. L. Wagner,¹⁹ J. Blouw,²⁰ J. L. Harton,²⁰ M. Krishnamurthy,²⁰ A. Soffer,²⁰ W. H. Toki,²⁰ R. J. Wilson,²⁰ J. Zhang,²⁰ T. Brandt,²¹ J. Brose,²¹ T. Colberg,²¹ M. Dickopp,²¹ R. S. Dubitzky,²¹ A. Hauke,²¹ E. Maly,²¹ R. Müller-Pfefferkorn,²¹ S. Otto,²¹ K. R. Schubert,²¹ R. Schwierz,²¹ B. Spaan,²¹ L. Wilden,²¹ D. Bernard,²² G. R. Bonneaud,²² F. Brochard,²² J. Cohen-Tanugi,²² S. Ferrag,²² E. Roussot,²² S. T'Jampens,²² Ch. Thiebaux,²² G. Vasileiadis,²² M. Verderi,²² A. Anjomshoa,²³ R. Bernet,²³ A. Khan,²³ D. Lavin,²³ F. Muheim,²³ S. Playfer,²³ J. E. Swain,²³ M. Falbo,²⁴ C. Borean,²⁵ C. Bozzi,²⁵ S. Dittongo,²⁵ L. Piemontese,²⁵ E. Treadwell,²⁶ F. Anulli,^{27,*} R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ D. Falciari,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,*} M. Piccolo,²⁷ Y. Xie,²⁷ A. Zallo,²⁷ S. Bagnasco,²⁸ A. Buzzo,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ F. C. Pastore,²⁸ C. Patrignani,²⁸ M. G. Pia,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ M. Morii,²⁹ R. Bartoldus,³⁰ R. Hamilton,³⁰ U. Mallik,³⁰ J. Cochran,³¹ H. B. Crawley,³¹ P.-A. Fischer,³¹ J. Lamsa,³¹ W. T. Meyer,³¹ E. I. Rosenberg,³¹ G. Grosdidier,³² C. Hast,³² A. Höcker,³² H. M. Lacker,³² S. Laplace,³² V. Lepeltier,³² A. M. Lutz,³² S. Plaszczynski,³² M. H. Schune,³² S. Trincaz-Duvoid,³² G. Wormser,³² R. M. Bionta,³³ V. Brigljević,³³ D. J. Lange,³³ M. Mugge,³³ K. van Bibber,³³ D. M. Wright,³³ M. Carroll,³⁴ J. R. Fry,³⁴ E. Gabathuler,³⁴ R. Gamet,³⁴ M. George,³⁴ M. Kay,³⁴ D. J. Payne,³⁴ R. J. Sloane,³⁴ C. Touramanis,³⁴ M. L. Aspinwall,³⁵ D. A. Bowerman,³⁵ P. D. Dauncey,³⁵ U. Egede,³⁵ I. Eschrich,³⁵ N. J. W. Gunawardane,³⁵ J. A. Nash,³⁵ P. Sanders,³⁵ D. Smith,³⁵ D. E. Azzopardi,³⁶ J. J. Back,³⁶ P. Dixon,³⁶ P. F. Harrison,³⁶ R. J. L. Potter,³⁶ H. W. Shorthouse,³⁶ P. Strother,³⁶ P. B. Vidal,³⁶ M. I. Williams,³⁶ G. Cowan,³⁷ S. George,³⁷ M. G. Green,³⁷ A. Kurup,³⁷ C. E. Marker,³⁷ P. McGrath,³⁷ T. R. McMahon,³⁷ S. Ricciardi,³⁷ F. Salvatore,³⁷ I. Scott,³⁷ G. Vaitsas,³⁷ D. Brown,³⁸ C. L. Davis,³⁸ J. Allison,³⁹ R. J. Barlow,³⁹ J. T. Boyd,³⁹ A. C. Forti,³⁹ J. Fullwood,³⁹ F. Jackson,³⁹ G. D. Lafferty,³⁹ N. Savvas,³⁹ E. T. Simopoulos,³⁹ J. H. Weatherall,³⁹ A. Farbin,⁴⁰ A. Jawahery,⁴⁰ V. Lillard,⁴⁰ J. Olsen,⁴⁰ D. A. Roberts,⁴⁰ J. R. Schieck,⁴⁰ G. Blaylock,⁴¹ C. Dallapiccola,⁴¹ K. T. Flood,⁴¹ S. S. Hertzbach,⁴¹ R. Kofler,⁴¹ V. G. Koptchev,⁴¹ T. B. Moore,⁴¹ H. Staengle,⁴¹ S. Willocq,⁴¹ B. Brau,⁴² R. Cowan,⁴² G. Sciolla,⁴² F. Taylor,⁴² R. K. Yamamoto,⁴² M. Milek,⁴³ P. M. Patel,⁴³ F. Palombo,⁴⁴ J. M. Bauer,⁴⁵ L. Cremaldi,⁴⁵ V. Eschenburg,⁴⁵ R. Kroeger,⁴⁵ J. Reidy,⁴⁵ D. A. Sanders,⁴⁵ D. J. Summers,⁴⁵ J. P. Martin,⁴⁶ J. Y. Nief,⁴⁶ R. Seitz,⁴⁶ P. Taras,⁴⁶ V. Zacek,⁴⁶ H. Nicholson,⁴⁷ C. S. Sutton,⁴⁷ N. Cavallo,^{48,†} G. De Nardo,⁴⁸ F. Fabozzi,⁴⁸ C. Gatto,⁴⁸ L. Lista,⁴⁸ P. Paolucci,⁴⁸ D. Piccolo,⁴⁸ C. Sciacca,⁴⁸ J. M. LoSecco,⁴⁹ J. R. G. Alsmiller,⁵⁰ T. A. Gabriel,⁵⁰ T. Handler,⁵⁰ J. Brau,⁵¹ R. Frey,⁵¹ M. Iwasaki,⁵¹ N. B. Sinev,⁵¹ D. Strom,⁵¹ F. Colecchia,⁵² F. Dal Corso,⁵² A. Dorigo,⁵² F. Galeazzi,⁵² M. Margoni,⁵² G. Michelon,⁵² M. Morandin,⁵² M. Posocco,⁵² M. Rotondo,⁵² F. Simonetto,⁵² R. Stroili,⁵² E. Torassa,⁵² C. Voci,⁵² M. Benayoun,⁵³ H. Briand,⁵³ J. Chauveau,⁵³ P. David,⁵³ Ch. de la Vaissière,⁵³ L. Del Buono,⁵³ O. Hamon,⁵³ F. Le Diberder,⁵³ Ph. Leruste,⁵³ L. Roos,⁵³ J. Stark,⁵³ S. Versillé,⁵³ P. F. Manfredi,⁵⁴ V. Re,⁵⁴ V. Speziali,⁵⁴ E. D. Frank,⁵⁵ L. Gladney,⁵⁵ Q. H. Guo,⁵⁵ J. Panetta,⁵⁵ C. Angelini,⁵⁶ G. Batignani,⁵⁶ S. Bettarini,⁵⁶ M. Bondioli,⁵⁶ M. Carpinelli,⁵⁶ F. Forti,⁵⁶ M. A. Giorgi,⁵⁶ A. Lusiani,⁵⁶ F. Martinez-Vidal,⁵⁶ M. Morganti,⁵⁶ N. Neri,⁵⁶ E. Paoloni,⁵⁶ M. Rama,⁵⁶ G. Rizzo,⁵⁶ F. Sandrelli,⁵⁶ G. Simi,⁵⁶ G. Triggiani,⁵⁶ J. Walsh,⁵⁶ M. Haire,⁵⁷ D. Judd,⁵⁷ K. Paick,⁵⁷ L. Turnbull,⁵⁷ D. E. Wagoner,⁵⁷ J. Albert,⁵⁸ P. Elmer,⁵⁸ C. Lu,⁵⁸ K. T. McDonald,⁵⁸ V. Miftakov,⁵⁸ S. F. Schaffner,⁵⁸ A. J. S. Smith,⁵⁸ A. Tumanov,⁵⁸ E. W. Varnes,⁵⁸ G. Cavoto,⁵⁹ D. del Re,⁵⁹ R. Faccini,⁵⁹ F. Ferrarotto,⁵⁹ F. Ferroni,⁵⁹ E. Lamanna,⁵⁹ E. Leonardi,⁵⁹ M. A. Mazzoni,⁵⁹ S. Morganti,⁵⁹ G. Piredda,⁵⁹ F. Safai Tehrani,⁵⁹ M. Serra,⁵⁹ C. Voena,⁵⁹ S. Christ,⁶⁰ R. Waldi,⁶⁰ T. Adye,⁶¹ B. Franek,⁶¹

N. I. Geddes,⁶¹ G. P. Gopal,⁶¹ S. M. Xella,⁶¹ R. Aleksan,⁶² G. De Domenico,⁶² S. Emery,⁶² A. Gaidot,⁶² S. F. Ganzhur,⁶² P.-F. Giraud,⁶² G. Hamel de Monchenault,⁶² W. Kozanecki,⁶² M. Langer,⁶² G. W. London,⁶² B. Mayer,⁶² B. Serfass,⁶² G. Vasseur,⁶² Ch. Yèche,⁶² M. Zito,⁶² N. Coptý,⁶³ M. V. Purohit,⁶³ H. Singh,⁶³ F. X. Yumiceva,⁶³ I. Adam,⁶⁴ P. L. Anthony,⁶⁴ D. Aston,⁶⁴ K. Baird,⁶⁴ N. Berger,⁶⁴ E. Bloom,⁶⁴ A. M. Boyarski,⁶⁴ F. Bulos,⁶⁴ G. Calderini,⁶⁴ M. R. Convery,⁶⁴ D. P. Coupal,⁶⁴ D. H. Coward,⁶⁴ J. Dorfan,⁶⁴ W. Dunwoodie,⁶⁴ R. C. Field,⁶⁴ T. Glanzman,⁶⁴ G. L. Godfrey,⁶⁴ S. J. Gowdy,⁶⁴ P. Grosso,⁶⁴ T. Haas,⁶⁴ T. Himel,⁶⁴ T. Hryn'ova,⁶⁴ M. E. Huffer,⁶⁴ W. R. Innes,⁶⁴ C. P. Jessop,⁶⁴ M. H. Kelsey,⁶⁴ P. Kim,⁶⁴ M. L. Kocian,⁶⁴ U. Langenegger,⁶⁴ D. W. G. S. Leith,⁶⁴ S. Luitz,⁶⁴ V. Luth,⁶⁴ H. L. Lynch,⁶⁴ H. Marsiske,⁶⁴ S. Menke,⁶⁴ R. Messner,⁶⁴ K. C. Moffeit,⁶⁴ R. Mount,⁶⁴ D. R. Muller,⁶⁴ C. P. O'Grady,⁶⁴ M. Perl,⁶⁴ S. Petrak,⁶⁴ H. Quinn,⁶⁴ B. N. Ratcliff,⁶⁴ S. H. Robertson,⁶⁴ L. S. Rochester,⁶⁴ A. Roodman,⁶⁴ T. Schietinger,⁶⁴ R. H. Schindler,⁶⁴ J. Schwiening,⁶⁴ V. V. Serbo,⁶⁴ A. Snyder,⁶⁴ A. Soha,⁶⁴ S. M. Spanier,⁶⁴ J. Stelzer,⁶⁴ D. Su,⁶⁴ M. K. Sullivan,⁶⁴ H. A. Tanaka,⁶⁴ J. Va'vra,⁶⁴ S. R. Wagner,⁶⁴ A. J. R. Weinstein,⁶⁴ W. J. Wisniewski,⁶⁴ D. H. Wright,⁶⁴ C. C. Young,⁶⁴ P. R. Burchat,⁶⁵ C. H. Cheng,⁶⁵ D. Kirkby,⁶⁵ T. I. Meyer,⁶⁵ C. Roat,⁶⁵ R. Henderson,⁶⁶ W. Bugg,⁶⁷ H. Cohn,⁶⁷ A. W. Weidemann,⁶⁷ J. M. Izen,⁶⁸ I. Kitayama,⁶⁸ X. C. Lou,⁶⁸ F. Bianchi,⁶⁹ M. Bona,⁶⁹ D. Gamba,⁶⁹ A. Smol,⁶⁹ L. Bosisio,⁷⁰ G. Della Ricca,⁷⁰ L. Lanceri,⁷⁰ P. Poropat,⁷⁰ G. Vuagnin,⁷⁰ R. S. Panvini,⁷¹ C. M. Brown,⁷² R. Kowalewski,⁷² J. M. Roney,⁷² H. R. Band,⁷³ E. Charles,⁷³ S. Dasu,⁷³ F. Di Lodovico,⁷³ A. M. Eichenbaum,⁷³ H. Hu,⁷³ J. R. Johnson,⁷³ R. Liu,⁷³ Y. Pan,⁷³ R. Prepost,⁷³ I. J. Scott,⁷³ S. J. Sekula,⁷³ J. H. von Wimmersperg-Toeller,⁷³ S. L. Wu,⁷³ Z. Yu,⁷³ T. M. B. Kordich,⁷⁴ and H. Neal⁷⁴

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720

⁶University of Birmingham, Birmingham B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697

¹³University of California at Los Angeles, Los Angeles, California 90024

¹⁴University of California at San Diego, La Jolla, California 92093

¹⁵University of California at Santa Barbara, Santa Barbara, California 93106

¹⁶University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064

¹⁷California Institute of Technology, Pasadena, California 91125

¹⁸University of Cincinnati, Cincinnati, Ohio 45221

¹⁹University of Colorado, Boulder, Colorado 80309

²⁰Colorado State University, Fort Collins, Colorado 80523

²¹Technische Universität Dresden, Institut für Kern und Teilchenphysik, D-01062, Dresden, Germany

²²Ecole Polytechnique, F-91128 Palaiseau, France

²³University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁴Elon University, Elon University, North Carolina 27244-2010

²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

²⁶Florida A&M University, Tallahassee, Florida 32307

²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

²⁹Harvard University, Cambridge, Massachusetts 02138

³⁰University of Iowa, Iowa City, Iowa 52242

³¹Iowa State University, Ames, Iowa 50011-3160

³²Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France

³³Lawrence Livermore National Laboratory, Livermore, California 94550

³⁴University of Liverpool, Liverpool L69 3BX, United Kingdom

³⁵University of London, Imperial College, London SW7 2BW, United Kingdom

³⁶Queen Mary, University of London, E1 4NS, United Kingdom

³⁷University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

³⁸University of Louisville, Louisville, Kentucky 40292

³⁹University of Manchester, Manchester M13 9PL, United Kingdom

⁴⁰University of Maryland, College Park, Maryland 20742

⁴¹University of Massachusetts, Amherst, Massachusetts 01003

⁴²Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139

⁴³McGill University, Montréal, Québec, Canada H3A 2T8

- ⁴⁴Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁵University of Mississippi, University, Mississippi 38677
⁴⁶Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Québec, Canada H3C 3J7
⁴⁷Mount Holyoke College, South Hadley, Massachusetts 01075
⁴⁸Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁴⁹University of Notre Dame, Notre Dame, Indiana 46556
⁵⁰Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831
⁵¹University of Oregon, Eugene, Oregon 97403
⁵²Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵³Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
⁵⁴Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
⁵⁵University of Pennsylvania, Philadelphia, Pennsylvania 19104
⁵⁶Università di Pisa, Scuola Normale Superiore and INFN, I-56010 Pisa, Italy
⁵⁷Prairie View A&M University, Prairie View, Texas 77446
⁵⁸Princeton University, Princeton, New Jersey 08544
⁵⁹Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶⁰Universität Rostock, D-18051 Rostock, Germany
⁶¹Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom
⁶²DAPNIA, Commissariat à l'Energie Atomique/Saclay, F-91191 Gif-sur-Yvette, France
⁶³University of South Carolina, Columbia, South Carolina 29208
⁶⁴Stanford Linear Accelerator Center, Stanford, California 94309
⁶⁵Stanford University, Stanford, California 94305-4060
⁶⁶TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
⁶⁷University of Tennessee, Knoxville, Tennessee 37996
⁶⁸University of Texas at Dallas, Richardson, Texas 75083
⁶⁹Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁰Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷¹Vanderbilt University, Nashville, Tennessee 37235
⁷²University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷³University of Wisconsin, Madison, Wisconsin 53706
⁷⁴Yale University, New Haven, Connecticut 06511
- (Received 7 September 2001; published 21 December 2001)

We measure the branching fractions of the $\psi(2S)$ meson to the leptonic final states e^+e^- and $\mu^+\mu^-$ relative to that for $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$. The method uses $\psi(2S)$ mesons produced in the decay of B mesons at the $Y(4S)$ resonance in a data sample collected with the BABAR detector at the Stanford Linear Accelerator Center. Using previous measurements for the $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ branching fraction, we determine the e^+e^- and $\mu^+\mu^-$ branching fractions to be $0.0078 \pm 0.0009 \pm 0.0008$ and $0.0067 \pm 0.0008 \pm 0.0007$, respectively.

DOI: 10.1103/PhysRevD.65.031101

PACS number(s): 13.20.Gd, 13.25.Gv, 14.40.Gx

The branching fraction of the $\psi(2S)$ to e^+e^- has previously been measured in e^+e^- collider experiments operating at the mass of the $\psi(2S)$ resonance [1] and in $p\bar{p}$ experiments [2,3]. The $\psi(2S) \rightarrow \mu^+\mu^-$ branching fraction has been measured with substantially larger uncertainty in e^+e^- experiments [4] and in π^-Be collisions [5]. This paper reports new measurements of these quantities by the BABAR experiment, operating at the PEP-II e^+e^- collider at the Stanford Linear Accelerator Center.

PEP-II collides 9 GeV electrons on 3.1 GeV positrons to create a center-of-mass system with an energy of 10.58 GeV moving along the z axis with a Lorentz boost of $\beta\gamma=0.56$. At this energy, $Y(4S)$ resonance production makes up 23% of the total hadronic cross section. The $Y(4S)$ is assumed to

decay 100% to a pair of B mesons. A large, clean sample of $\psi(2S)$ mesons is produced in the B decays. The e^+e^- and $\mu^+\mu^-$ branching fractions are obtained through their ratio to $J/\psi \pi^+ \pi^-$, which is known with much better precision. This technique provides a significantly lower uncertainty on the $\mu^+\mu^-$ branching fraction than the current world average.

The data set used for this analysis corresponds to an integrated luminosity of $20.33 \pm 0.30 \text{ fb}^{-1}$ recorded at 10.58 GeV, and contains $(22.3 \pm 0.4) \times 10^6$ $Y(4S)$ mesons. An additional 2.6 fb^{-1} has been recorded at an energy 40 MeV below the $Y(4S)$ resonance.

The BABAR detector is described in detail in Ref. [6]. The momenta of charged particles are measured and their trajectories reconstructed with two detector systems located in a 1.5 T solenoidal magnetic field: a five-layer, double-sided silicon vertex tracker (SVT) and a 40 layer drift chamber (DCH). The fiducial volume covers the polar angular region $0.41 < \theta < 2.54$ rad, which is 86% of the solid angle in

*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

the center of mass. The transverse momentum resolution is 0.47% at 1 GeV/ c .

The energies of electrons and photons are accurately measured by a CsI(Tl) calorimeter (EMC) in the fiducial volume $0.41 < \theta < 2.41$ rad (84% of the center-of-mass solid angle) with an energy resolution at 1 GeV of 3.0%. Muons are detected in the IFR—the flux return of the solenoid, which is instrumented with resistive plate chambers. The DIRC, a unique Cherenkov radiation detection device, identifies charged particles.

The branching fractions of interest are obtained by comparison to that of $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$. The number of $\psi(2S)$ mesons reconstructed in the final states e^+e^- (N_{ee}), $\mu^+\mu^-$ ($N_{\mu\mu}$) and $J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow e^+e^-$ ($N_{ee\pi\pi}$) or $J/\psi \rightarrow \mu^+\mu^-$ ($N_{\mu\mu\pi\pi}$), is related to the total number of $\psi(2S)$ mesons produced in our data set $N_{\psi(2S)}$ by

$$N_{ee} = N_{\psi(2S)} \cdot \mathcal{B}_{ee} \cdot \epsilon_{ee}, \quad (1)$$

$$N_{\mu\mu} = N_{\psi(2S)} \cdot \mathcal{B}_{\mu\mu} \cdot \epsilon_{\mu\mu}, \quad (2)$$

$$N_{ee\pi\pi} = N_{\psi(2S)} \cdot \mathcal{B}_{J/\psi \pi^+ \pi^-} \cdot \mathcal{B}_{J/\psi \rightarrow ee} \cdot \epsilon_{ee\pi\pi}, \quad (3)$$

$$N_{\mu\mu\pi\pi} = N_{\psi(2S)} \cdot \mathcal{B}_{J/\psi \pi^+ \pi^-} \cdot \mathcal{B}_{J/\psi \rightarrow \mu\mu} \cdot \epsilon_{\mu\mu\pi\pi}. \quad (4)$$

\mathcal{B}_{ee} , $\mathcal{B}_{\mu\mu}$, and $\mathcal{B}_{J/\psi \pi^+ \pi^-}$ are the branching fractions of the $\psi(2S)$ to e^+e^- , $\mu^+\mu^-$, and $J/\psi \pi^+ \pi^-$, respectively. We use world averages for $\mathcal{B}_{J/\psi \rightarrow ee}$, the J/ψ branching fraction to e^+e^- , and for $\mathcal{B}_{J/\psi \rightarrow \mu\mu}$, the branching fraction to $\mu^+\mu^-$ [7]. ϵ_{ee} and $\epsilon_{\mu\mu}$ are the efficiencies for events containing $\psi(2S)$ mesons decaying to e^+e^- and $\mu^+\mu^-$ respectively to satisfy the event selection and meson reconstruction requirements; $\epsilon_{ee\pi\pi}$ and $\epsilon_{\mu\mu\pi\pi}$ are the efficiencies for $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ decays with $J/\psi \rightarrow e^+e^-$ and $J/\psi \rightarrow \mu^+\mu^-$, respectively.

Equations (1), (3), and (4) can be combined to give two expressions for the e^+e^- to $J/\psi \pi^+ \pi^-$ branching ratio:

$$\frac{\mathcal{B}_{ee}}{\mathcal{B}_{J/\psi \pi^+ \pi^-}} = \mathcal{B}_{J/\psi \rightarrow ee} \cdot \frac{N_{ee}}{N_{ee\pi\pi}} \cdot \frac{\epsilon_{ee\pi\pi}}{\epsilon_{ee}}, \quad (5)$$

$$\frac{\mathcal{B}_{ee}}{\mathcal{B}_{J/\psi \pi^+ \pi^-}} = \mathcal{B}_{J/\psi \rightarrow \mu\mu} \cdot \frac{N_{ee}}{N_{\mu\mu\pi\pi}} \cdot \frac{\epsilon_{\mu\mu\pi\pi}}{\epsilon_{ee}}. \quad (6)$$

Similarly,

$$\frac{\mathcal{B}_{\mu\mu}}{\mathcal{B}_{J/\psi \pi^+ \pi^-}} = \mathcal{B}_{J/\psi \rightarrow ee} \cdot \frac{N_{\mu\mu}}{N_{ee\pi\pi}} \cdot \frac{\epsilon_{ee\pi\pi}}{\epsilon_{\mu\mu}}, \quad (7)$$

$$\frac{\mathcal{B}_{\mu\mu}}{\mathcal{B}_{J/\psi \pi^+ \pi^-}} = \mathcal{B}_{J/\psi \rightarrow \mu\mu} \cdot \frac{N_{\mu\mu}}{N_{\mu\mu\pi\pi}} \cdot \frac{\epsilon_{\mu\mu\pi\pi}}{\epsilon_{\mu\mu}}. \quad (8)$$

A number of systematic errors due to uncertainties in efficiency cancel in these expressions.

We obtain a $B\bar{B}$ enriched sample by requiring events to have visible energy E greater than 4.5 GeV and a ratio of the second to the zeroth Fox-Wolfram moment, R_2 [8], less than 0.5. Both E and R_2 are calculated from tracks and neutral clusters in the respective fiducial volumes noted above. The

same tracks are used to construct a primary event vertex, which is required to be located within 6 cm of the beam spot in z and within 0.5 cm of the beam line. The beam spot rms size is approximately 0.9 cm in z , 120 μm horizontally, and 5.6 μm vertically.

There must be at least three tracks in the fiducial volume satisfying the following quality criteria: they must have transverse momentum greater than 0.1 GeV/ c , momentum less than 10 GeV/ c , at least 12 hits in the DCH, and approach within 10 cm of the beam spot in z and within 1.5 cm of the beam line.

Finally, to suppress a substantial background from radiative Bhabha ($e^+e^- \gamma$) events in which the photon converts to an e^+e^- pair, five or more tracks are required in events containing $\psi(2S) \rightarrow e^+e^-$ or $J/\psi \rightarrow e^+e^-$ candidates.

The efficiency of the event selection—and the meson reconstruction efficiency described below—is calculated with a complete detector simulation of $B \rightarrow \psi(2S)X$ events [9]. The simulation of $\psi(2S)$ and J/ψ decays to lepton pairs includes final state radiation [10]. The event selection efficiencies are 0.912 ± 0.002 for $\psi(2S) \rightarrow e^+e^-$, 0.945 ± 0.002 for $\psi(2S) \rightarrow \mu^+\mu^-$, 0.967 ± 0.001 for $e^+e^- \pi^+ \pi^-$, and 0.972 ± 0.001 for $\mu^+\mu^- \pi^+ \pi^-$. The difference in the e^+e^- and $\mu^+\mu^-$ efficiencies is due largely to the requirement of five tracks. The quoted uncertainties are those due to simulation statistics only. The event efficiencies appear as ratios in Eqs. (5)–(8); the systematic errors on the ratios are small compared to the other uncertainties and systematic errors discussed below.

The lepton candidates used to construct J/ψ or $\psi(2S)$ mesons via e^+e^- or $\mu^+\mu^-$ decays must be in the restricted angular region $0.41 < \theta < 2.41$ rad and satisfy the track quality criteria listed above.

Electron candidates must include an energy deposition in the EMC of at least three crystals, with shape consistent with an electromagnetic shower and magnitude at least 75% of the track momentum. At least one candidate must have energy between 89% and 120% of the track momentum and a Cherenkov signal in the DIRC consistent with the expectation for an electron. If possible, photons radiated by electrons traversing material prior to the DCH are recombined with the track. Such photons must have EMC energy greater than 30 MeV, a polar angle θ within 35 mrad of the electron direction and an azimuth that is either within 50 mrad of the electron direction or between the electron direction and the location of the electron shower in the EMC.

Muon candidates must deposit less than 0.5 GeV in the EMC (2.3 times the minimum-ionizing peak), penetrate at least two interaction lengths λ of material, and have a pattern of hits consistent with the trajectory of a muon. We require the material traversed by one candidate be within 1λ of that expected for a muon; for the other candidate, this is relaxed to 2λ .

The J/ψ or $\psi(2S)$ meson mass is obtained in an l^+l^- final state after constraining the two tracks to a common origin.

The reconstruction of $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ uses a $J/\psi \rightarrow e^+e^-$ candidate with mass between 3.05 and 3.12 GeV/ c^2 or a $J/\psi \rightarrow \mu^+\mu^-$ candidate with $3.07 < m < 3.12$ GeV/ c^2 .

74% of $J/\psi \rightarrow e^+e^-$ decays and 91% of $J/\psi \rightarrow \mu^+\mu^-$ fall within these ranges. All tracks in the fiducial volume not used in the J/ψ reconstruction are used as pion candidates. To avoid systematic errors and retain high efficiency, the tracks are not required to satisfy any specific quality requirements. A pair of oppositely-charged pions is required to have mass $m_{\pi\pi}$ in the region $0.45 < m_{\pi\pi} < 0.60 \text{ GeV}/c^2$. The $\psi(2S)$ mass is obtained after constraining the four tracks in the final state to a common origin.

$\psi(2S)$ candidates in all final states are required to have momentum less than $1.6 \text{ GeV}/c$ as measured in the $Y(4S)$ rest frame. This requirement is fully efficient for $\psi(2S)$ mesons produced in B decays.

The J/ψ and $\psi(2S)$ reconstruction efficiencies are determined by simulation and include contributions from acceptance, track quality, particle identification and, for $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, the J/ψ and $\pi^+ \pi^-$ mass windows. The efficiency and systematic error on lepton identification have been obtained from data by comparing the ratio of J/ψ mesons in B decays in which one or both leptons satisfy the requirements. The efficiency and systematic error of the track-quality selection have been studied by comparing the independent SVT and DCH tracking efficiencies in hadronic events. The meson reconstruction efficiency is 0.602 ± 0.004 for the e^+e^- case, 0.535 ± 0.004 for $\mu^+\mu^-$, 0.207 ± 0.002 for $e^+e^- \pi^+ \pi^-$, and 0.211 ± 0.002 for $\mu^+\mu^- \pi^+ \pi^-$, where the uncertainties are simulation statistics only.

The e^+e^- efficiency is higher than $\mu^+\mu^-$ in $\psi(2S) \rightarrow l^+l^-$ or $J/\psi \rightarrow l^+l^-$ reconstruction because electron identification is more efficient than muon identification. Conversely, a J/ψ decaying to e^+e^- is less likely to be reconstructed in the specified mass window than one decaying to $\mu^+\mu^-$. Together, these two effects result in little difference between the $e^+e^- \pi^+ \pi^-$ and $\mu^+\mu^- \pi^+ \pi^-$ efficiencies. Overall, the $J/\psi \pi^+ \pi^-$ efficiencies are lower than l^+l^- due to the reconstruction of the pion pair. The efficiencies appearing in Eqs. (1)–(4) are the product of these meson reconstruction efficiencies and the event selection values given earlier.

Lepton identification uncertainty is 1.8% for e^+e^- and 1.4% for $\mu^+\mu^-$, and cancels in branching ratios where the $\psi(2S)$ and J/ψ decay to the same final state, Eqs. (5) and (8). A 2.4% systematic error on the efficiency of the track quality requirements applied to the J/ψ and $\psi(2S)$ in the l^+l^- final state cancels in all four ratios.

The number of mesons in the e^+e^- and $\mu^+\mu^-$ final states is extracted by a fit to the mass distribution of candidates (Fig. 1). A third-order Chebychev polynomial is used for backgrounds. The signals are fit by probability distribution functions (pdf's) obtained from a complete simulation of $B \rightarrow \psi(2S)X$ events, with $\psi(2S) \rightarrow e^+e^-$ or $\psi(2S) \rightarrow \mu^+\mu^-$. Only candidates constructed from the correct combination of particles are used in the pdf. The signal pdf's are convoluted with a Gaussian distribution to match the mass resolution of $12 \text{ MeV}/c^2$ observed in a data sample of 14 000 $J/\psi \rightarrow \mu^+\mu^-$ decays.




FIG. 1. Mass distribution of (a) $\psi(2S) \rightarrow e^+e^-$ and (b) $\psi(2S) \rightarrow \mu^+\mu^-$ candidates.

Despite the algorithm to recover radiated photons, the pdf for the e^+e^- final state is sensitive to the fraction of events in which one or both electrons undergo bremsstrahlung. The pdf is adjusted to reflect the fraction obtained in a study of the mass distribution of 15 000 $J/\psi \rightarrow e^+e^-$ decays in data. To enhance the sensitivity of the study, the algorithm to recover radiated photons is not used in the reconstruction of the J/ψ .

For $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, an analogous fit procedure is performed to the distribution of the mass difference between the $\psi(2S)$ and the J/ψ candidates (Fig. 2). This quantity reduces the impact of J/ψ mass resolution, including final state radiation and bremsstrahlung. The distribution predicted by the simulation is convoluted with a Gaussian distribution whose standard deviation is left as a free parameter in the fit. The mass difference resolution is $3.2 \text{ MeV}/c^2$.

The signal yields returned by the fits are 552 ± 50 for e^+e^- , 437 ± 44 for $\mu^+\mu^-$, 474 ± 44 for $e^+e^- \pi^+ \pi^-$, and 498 ± 42 for $\mu^+\mu^- \pi^+ \pi^-$, where errors are statistical only. Systematic errors on the fitting technique are obtained by performing the fits on multiple simulated data sets containing both signal and background events. Additional contributions come from varying the mass regions included in the fit and increasing or decreasing the power of the background polynomial. Fitting systematics are 2.3% for e^+e^- , 5.3% for $\mu^+\mu^-$, 5.4% for $e^+e^- \pi^+ \pi^-$, and 2.1% for $\mu^+\mu^- \pi^+ \pi^-$. These systematic errors are conservative in the sense that the procedure to derive them incorporates a component of the statistical error, which would be reduced with additional data.

We repeat the analysis with the data recorded below the $Y(4S)$ resonance. The total $\psi(2S)$ yield, summed over the four modes, is 5 ± 12 events, indicating that the contribution




FIG. 2. Mass difference between the $\psi(2S)$ and J/ψ candidates in the decay $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ with the J/ψ reconstructed in the (a) e^+e^- and (b) $\mu^+\mu^-$ final states.

of continuum-produced $\psi(2S)$ mesons is negligible in the on-resonance sample.

The two values for the e^+e^- to $J/\psi\pi^+\pi^-$ branching ratio obtained with Eqs. (5) and (6) are in good agreement: the result found with $\mu^+\mu^-\pi^+\pi^-$ is 0.97 ± 0.14 times that with $e^+e^-\pi^+\pi^-$. By construction, this ratio is identical for the $\mu^+\mu^-$ final state. The results from Eqs. (5) and (6) are combined, distinguishing correlated and uncorrelated statistical and systematic errors, to give

$$\mathcal{B}_{ee}/\mathcal{B}_{J/\psi\pi^+\pi^-} = 0.0252 \pm 0.0028 \pm 0.0011, \quad (9)$$

where the first error is statistical and the second systematic. Similarly, Eqs. (7) and (8) are combined to obtain

$$\mathcal{B}_{\mu\mu}/\mathcal{B}_{J/\psi\pi^+\pi^-} = 0.0216 \pm 0.0026 \pm 0.0014. \quad (10)$$

The systematic errors are dominated by the fitting technique. Other contributions, which are the same for both results, include 1.6% for particle identification, 1.2% for the uncertainty in J/ψ branching fractions, and 0.9% for differences between the simulated and measured [11] $\pi^+\pi^-$ mass and angular distributions in the $J/\psi\pi^+\pi^-$ final states.

We use the current world average value of 0.310 ± 0.028 for the $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ branching fraction [7] to extract results for the $\psi(2S)$ leptonic branching fractions:

$$\mathcal{B}_{ee} = 0.0078 \pm 0.0009 \pm 0.0008, \quad (11)$$

$$\mathcal{B}_{\mu\mu} = 0.0067 \pm 0.0008 \pm 0.0007. \quad (12)$$

The ratio of the leptonic branching fractions can be derived without the use of the $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ sample:

$$\frac{\mathcal{B}_{\mu\mu}}{\mathcal{B}_{ee}} = \frac{N_{\mu\mu}}{N_{ee}} \cdot \frac{\epsilon_{ee}}{\epsilon_{\mu\mu}} = 0.86 \pm 0.12 \pm 0.05. \quad (13)$$

The systematic error is dominated by the uncertainty in the fitting technique.

In summary, we have measured the branching ratios $\mathcal{B}_{ee}/\mathcal{B}_{J/\psi\pi^+\pi^-}$ and $\mathcal{B}_{\mu\mu}/\mathcal{B}_{J/\psi\pi^+\pi^-}$. We multiply these by the world average for the $J/\psi\pi^+\pi^-$ branching fraction to obtain the branching fraction of the $\psi(2S)$ to e^+e^- and to $\mu^+\mu^-$. These results are consistent with earlier measurements, but have, in the case of $\mu^+\mu^-$, a substantially smaller uncertainty.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF (Germany), INFN (Italy), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the Swiss NSF, A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

[1] G. J. Feldman and M. L. Perl, Phys. Rep. **33**, 285 (1977).
 [2] Fermilab E760 Collaboration, T. A. Armstrong *et al.*, Phys. Rev. D **55**, 1153 (1997).
 [3] Fermilab E835 Collaboration, M. Ambrogiani, *et al.*, Phys. Rev. D **62**, 032004 (2000).
 [4] E. Hilger *et al.*, Phys. Rev. Lett. **35**, 625 (1975).
 [5] E672 and E706 Collaborations, A. Gribushin *et al.*, Phys. Rev. D **53**, 4723 (1996).
 [6] BABAR Collaboration, B. Aubert *et al.*, SLAC-PUB-8569, hep-ex/0105044.

[7] Particle Data Group, D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
 [8] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
 [9] "GEANT Detector Description and Simulation Tool," Version 3.21, CERN Program Library Long Writeup W5013, 1994.
 [10] E. Barberio, B. van Eijk, and Z. Was, Comput. Phys. Commun. **66**, 115 (1991).
 [11] BES Collaboration, J. Z. Bai *et al.*, Phys. Rev. D **62**, 032002 (2000).